

COMPANY

**PRODUCT**

PROFILE



- ◆ INSECTICIDE
- ◆ FUNGICIDE
- ◆ HERBICIDE
- ◆ PGR



**SHIVCHEM**  
AGRO PVT. LTD.



## About Company:

**SHIVCHEM AGRO PVT. LTD.** We are committed to educate the farmer to use crop and pest specific crop protection products judiciously. We are working towards adding new chemistry in the product basket to ensure safer products for users. Our sales and marketing team strategically guides the farmers with the adequate and newly adopted techniques which range from teaching sowing pattern, harvesting pattern and shifting from one crop to another in reference to ongoing weather shifts. In the first phase, we opened our operations in North India and are steadily shifting towards Central India. In a little span of time, our operations will be functional throughout the country.

In our operational states such as Punjab, Uttarparadesh, Haryana, Karnataka, A.P., M.P., Telangana & Rajasthan, we have a presence across 500 distributors and 2000 retailers. Our brand is sharing shelves with top leading brands among their categories. Our quality products and their attractive packaging is generating repeated purchase from the farmers which is further building their trust on our company and organisation as a whole. We have experienced magnificent growth over two years since our inception and by 2020 we are determined to reach higher limits.



## Mission :

Educate, Evolve and Enhance - Strategy of Shivchem Agro Pvt. Ltd.

## Vision :

To emerge as a leading corporate in delivering agro-solutions and imparting knowledge to end user alongside strengthening manufacturing capabilities, strategic partnerships to superior technology and nurturing innovation across the value chain.

## Values :

Trust / Equality / Partnering / Safety / Care



## INSECTICIDE



### KRIDAN 4G

Cartap Hydrochloride 4% GR

Insecticide

**KRIDAN 4G** is a Broad spectrum systemic insecticide with contact and stomach action for Soil Application.

**Crop :** Paddy.

**Applied on :** Whorl Maggot, Sucking & Chewing Pests, Stem Borers, Leaf folder, etc

**Dosage Per Ha\*:** 18-25 Kg per HA

**Packing :** 5 kg, 1 kg.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Carbofuron 3% CG

Insecticide

**CARBAIN 3G** is a systemic Nematicide control all types of insects

**Crop:** Sugarcane, cotton, Paddy, chilli.

**Applied on :** Thrips, Aphids, Jassids, Nematicide.

**Dosage :** 25-33kg per Ha

**Packing :** 5 kg, 1 kg.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### CARBAIN 3G



### E-MAX

Emamectin Benzoate 5% SG

Insecticide

**E-MAX** is biological Insecticide/acricide derived having stomach action

**Crop :** Cotton, Okra

**Applied on :** Bollworms, Fruits & shoot borer

**Dose :** 220 gms per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.





## SHIVGENT

Fipronil 0.3% GR

Insecticide

**SHIVGENT** is a broad spectrum moderately systemic insecticide, toxic by contact and ingestion.

**Crop :** Paddy, Sugarcane

**Applied on :** Stem, Root, Shoot borers, Whorl maggot, rice gall midg, etc

**Dosage Per Ha\* :** 25-33 Kg/H/A

**Packing :** 5 kg, 1 kg.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## THIA-KING

Insecticide

Thiamethoxam 30% FS

**THIA-KING** is provides of a wide range of early season sucking and chewing pests, It is used to Treat the seeds.

**Crop :** Cotton, Wheat, Soybeans, Corn, Sorghum.

**Applied on :** Aphids, Wireworms, Flea beetles, Lea miners.

**Dosage Per Ha\* :** 10-20 ml water per kg seeds

**Packing :** 1 Ltr, 500 ml. 250 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## CHLOMATE 10G

Chlorpyrifos 10% GR

Insecticide

**CHLOMATE 10 G** is a insecticides. It is used for controlling of chewing and sucking insects in various crops.

**Crop :** Sugarcane, Paddy

**Applied on :** Stem borer, leaf folder, whorl maggot, shoot borer & root borer.

**Dosage Per Ha\*:** 7.5 to 10 kg / acre

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## INSECTICIDE



### KILLBAN-50

Chlorpyrifos 50% EC

Insecticide

**KILLBAN-50** is an Organophosphorous Broad Spectrum Non Systemic insecticide with contact, stomach and vapour action.

**Crop :** Paddy, Cotton, Termite control Pre & Post building construction

**Applied on :** Termites, Stem borer, Leaf roller, Bollworm, etc.

**Dosage Per Ha\*:** 1000-1200 ML per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### MAXO STAR

Thiamethoxam 25% WG Broad Spectrum systemic Insecticide

**MAXO STAR** is a Broad Spectrum Systemic insecticide with Contact and Stomach action

**Crop:** Paddy, Soyabean, Cotton, Vegetables.

**Applied on :** Stem borer, Leaf borer, BPH, Gall midge Insects, etc.

**Dosage :** 100 gm per HA in 500 - 700 litre water

**Packing :** 1 kg, 500 g, 250 g, 100 g.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### KILLBAN-20

Chlorpyrifos 20% EC

Insecticide

**KILLBAN-20** is an Organophosphorous Broad Spectrum Non Systemic insecticide with contact, stomach and vapour action.

**Crop :** Paddy, Gram, Cotton, Sugarcane, Ground Nut, Apple, Citrus

**Applied on :** Leaf roller, Aphid, Termites, Hairy Caterpillar, Stem borer, BPH, White Fly, etc

**Dosage Per Ha\*:** 500 - 1000 ML per HA in 500 - 700 litre water

**Packing :** 5 Ltr, 1 Ltr, 500ml, 250ml, 100ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## INSECTICIDE



### ARGON 505

Chlorpyrifos 50% + Cypermethrin 5% EC Insecticide

**ARGON 505** is broad spectrum Insectide.

**Crop :** Cotton

**Applied on :** Aphid, Jassid, Thrips, Pink bollworm & White Fly.

**Dose :** 1 Litre per HA in 500 - 700 Litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### APOCYPER-25

Cypermethrin 25% EC

Insecticide

**APOCYPER-25** is a synthetic pyrethroid Non Systemic insecticide with contact and stomach action.

**Crop :** Cotton, Vegetables, etc.

**Applied on :** Bollworm, Jassid, Thrip, Shoot & Fruit borer, etc.

**Dosage :** 250-400 ml per HA in 500-700 Ltr. of water.

**Packing :** 5 Ltr, 1 Ltr, 500ml, 250ml, 100ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### APOCYPER-10

Cypermethrin 10% EC

Insecticide

**APOCYPER-10** is a synthetic pyrethroid Non Systemic insecticide with contact and stomach action.

**Crop :** Cotton, Vegetables, etc.

**Applied on :** Bollworm, Jassid, Thrip, Shoot & Fruit borer, etc.

**Dosage :** 250 ml per HA in 500-700 Ltr. of water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



INSECTICIDE



## IMAAN

Imidacloprid 17.8% SL

Insecticide

**IMAAN** is a Broad Spectrum systemic and contact action insecticide.

**Crop:** Cotton, Sugarcane, Paddy, Okara, Sunflower

**Applied on:** Aphid, White Fly, Jassid, Termite, Thrips

**Dosage Per Ha\*:** 125-250 ML in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Insecticide

## ALPHAMEEN

Alphacyperemthrin 10% EC

**ALPHAMEEN** is highly active broad spectrum Insecticide having both contact & stomach action.

**Crop:** Cotton. Vegetables.

**Applied on:** Bollworm .

**Dosage Per Ha\*:** 165-280 ML per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## PROFOUND SUPER

Profenofos 40% + Cypermethrin 4% EC

Insecticide

**PROFOUND SUPER** is a Broad spectrum systemic insecticide with contact and stomach action.

**Crop:** Cotton. Chilli

**Applied on :** Boll worms Insects.

**Dosage Per Ha\*:** 1000-1500 ML per HA in 500 - 700 litre water

**Packing :** 1 Ltr, 500ml, 250ml, 100ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## INSECTICIDE



### REGAL

Fipronil 5% SC

Insecticide

**REGAL** is systemic and contact Insecticide

**Crop :** Sugarcane, Rice, Cabbage & Chilli

**Applied on :** Early shoot borer, stem borer, leaf folder, aphid, Thrips

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

**Dose :** 1-1.5 litre per HA in 500 - 700 litre water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### HIGH STAR

Insecticide

Bifenthrin 10% EC

**High Star** is broad spectrum Insecticide having Contact & Stomach action.

**Crop :** Cotton, Sugarcane, Rice

**Applied on :** Bollworm, White Fly, Termite, Stem borer, Leaf Folder

**Dosege :** 800-1000 ML per HA in 500 - 700 litre water

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### DECAFOL

Dicofol 18.5% EC

INSECTICIDE

**Decofol** is controlling many species of phytophagous mite on a wide range of crops, including fruits, vines, ornamental plants and vegetables.

**Crop :** Ground Nut, Okra, Sorghum, Tea

**Dosage Per Ha\*:** 1 Ltr. per HA in 400 - 1200 ml litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## INSECTICIDE



### PRO FOUND

Profenofos 50% EC

Insecticide

**Pro Found** is a non-systemic insecticide, acaricide with contact and stomach poison. Exhibits a translaminar effect, quick knockdown action with prolonged residual activity. Controls of insects (particularly Lepidoptera) and mites on cotton, maize, sugar beet, soya beans, potatoes, vegetables, tobacco and other crops.

**Dosage :** 1000 - 2000 ml per HA in 500 - 1000 litres of water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### CHECK MATE

Pymetrozine 50% WG

Insecticide

**Check Mate** has the best known chemistry for effective control on BPH, It is a systemic and trans laminar insecticide which paralyzes the hopper, stops egg-laying and the insects get killed from starvation.

**Crop :** Paddy

**Applied on :** Brown Plant Hopper

**Packing :** 1 kg, 500 g

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### INDOVANT

Indoxacarb 14.5 % SC

Insecticide

**INDOVANT** is a broad spectrum contact and stomach action insecticide.

**Crop :** Cotton.

**Applied on :** American Boll worm.

**Dosage :** 500 ml per HA in 500-700 Ltr. of water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## AKALUX

Quinalphos 25% EC

INSECTICIDE

**AKALUX** is an Organophosphorous broad Spectrum insecticide with contact and stomach action & quick knock down effect.

**Crop :** Paddy, wheat, Red Gram, Chillies, Mustard, Cotton, Etc.

**Applied on :** Hispa, Pod Borer, Scales, Thrips, brown plant hopper, Leaf roller, Stem borer, etc

**Dosage :** 1-1.5 litre per HA in 500 - 700 litre water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Insecticide

## AMITHION

Malathion 50% EC

**Heriattra** is a selective systemic pre and post emergence herbicide used for control of annual broad leaved weeds and grasses.

**Crop :** Maize, Bajra, Sugarcane, Potato, etc

**Dosage Per Ha\*:** 1-2 Kg per HA in 500 - 700 litre water

**Packing :** 500 gm.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## RIGOR

Dimethoate 30% EC

INSECTICIDE

**RIGOR** is a broad spectrum insecticide.

**Crop :** Mango, Potato, Banana, Cabbage, Brinjal, Okra, Chilli, Tomato, Onion.

**Fungal Diseases :** Aphid, Leaf hopper, White fly, Stem borer, Thrips, Mite etc

**Dosage Per Ha\*:** 1 Ltr. per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



INSECTICIDE



## LAMBTHRIN-5

Lambdacyhalothrin 5% EC

INSECTICIDE

**Lambthrin - 5** is a contact and systemic insecticide.

**Crop :** Cotton

**Applied on :** Bollworm, Jassids, Thrips, Fruit/ Shoot Borer etc

**Dosage Per Ha\*:** 150 - 250 ML in 500 - 700 litre of water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## LAMBTHRIN-2.5

Lambdacyhalothrin 2.5% EC

INSECTICIDE

**Lambthrin 2.5** is a contact and systemic insecticide.

**Crop :** Cotton

**Applied on :** Bollworm, Jassids, Thrips, Fruit/ Shoot Borer etc

**Dosage Per Ha\*:** 300 - 500 ML in 500 - 700 litre of water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## LAMBTHRIN

Lambda Cyhalothrin 4.9% CS

Insecticide

**Lambthrin** is an Insecticide in the form of Capsule suspension formulation.

**Crop :** Cotton, Paddy, Tomato, Okara, Brinjal

**Applied on :** Bollworm, Leaf Folder, Fruit borer, Thrips

**Dose :** 250-500 - 700 ML per HA in 500 - 700 Litre water

**Packing :** 1 Ltr, 500ml, 250ml, 100ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

INSECTICIDE



## IMAAN-G

Imidacloprid 0.3% GR

INSECTICIDE

**IMAAN-** is a systemic soil application insecticide containing 0.3% GR Imidacloprid active ingredient. Imidacloprid is used to control stem borer in paddy.

**Crop :** Paddy,

**Applied on :** Stem borer

**Packing :** 2.5 kg.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## IMAAN 30.5

INSECTICIDE

Imidacloprid 30.5% SC

**Imaan 30.5** is systemic and contact Insecticide

**Crop :** Rice, Cotton

**Applied on :** Brown Plant hopper, Aphid, Jassid, Thrips.

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## NARUTO

Novaluron 10% EC

Insecticide

Naruto is Insecticides are substances used to kill insects. They include ovicides and larvicides used against insect eggs and larvae, respectively. Insecticides are used in agriculture, medicine, industry, and consumer

**Crop :** Cotton, Cabbage, Tomato

**Applied on :** American bollworm, fruit-pod borer, diamondback motti, tobacco caterpillar

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## INSECTICIDE



### KRITAP-50

Cartap Hydrochloride 50% SP

INSECTICIDE

**Kritap-50** is a Broad spectrum Systemic insecticide with contact and stomach action.

**Crop :** Paddy.

**Applied on :** Sucking Stem borer, Leaf Folder & Chewing Pests

**Packing :** 1 kg, 500 g

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Emamectin Benzoate 1.9% EC

### EMAX-L

INSECTICIDE

**German Super** is systemic and contact Insecticide

**Crop :** Rice, Cotton

**Applied on :** Brown Plant hopper, Aphid, Jassid, Thrips.

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### RATOFF

Zinc Phosphide 80% w/w

Rodeniticde

**Ratoff** is Zinc phosphide is only used as a rodenticide. It is made into bait that will attract the pest, such as gophers, ground squirrels, or field mice.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## HERBICIDE



### KAKASHI

Quizalofop Ethyl 10% EC

Herbicide

**Kakashi** is a selective systemic herbicide, has a high selectivity between grass weeds & dicotyledons crops.

**Crop** : Soybean.

**Applied on** : Love Grass (Eragrostis Ipilosa), Crab Grass (Digitaria Sanguinalis) Wild Finger/ Makra Grass, Viper Grass, Barnyard Grass, Sanwa/Samel, Brown Top Millet

**Packing** : 1 Ltr, 500 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### GRIPPER

Metsulfuron Methyl 20% WP

Herbicide

**Gripper** is a selective post emergence herbicide with broad spectrum activity. Used for one shot control of wide range of broad leaf weeds.

**Crop**: Generally for Wheat

**Applied on**: Melilotus Indica, Melilotus alba, Vicia Satia, etc

**Dosage Per Ha\***: 8 gm per Acre

**Packing** : 8 gm + 200 ml Surfactant (1 Unit)

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### WEEDCHLOR

Butachlor 50% EC

Herbicide

**Azura-58** is non selective, contact, post emergent herbicide.

**Crop** : Potato, Cotton, Rubber, Wheat, Rice, Apple, Grapes

**Packing** : 1 Litre, 400 ML

**Dosage Per HA\*** : 1.25-3 Litre per HA in 500-700 litre Water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

HERBICIDE



## GRASS OUT

Glyphosate 41% SL

Herbicide

**GRASS OUT** is a non selective post emergent herbicide which is effective against annual and perennial grasses.

**Crop :** Tea, Non Cropped areas, Etc.

**Dosage Per Ha\*:** 2-3 litre per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## GLYWELL

Glyphosate 41% SL


Herbicide

**GRASS OUT** is a non selective post emergent herbicide which is effective against annual and perennial grasses.

**Crop :** Tea, Non Cropped areas, Etc.

**Dosage Per Ha\*:** 2-3 litre per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## GLYWELL-71

Ammonium Salt of Glyphosate 71% SG Systemic Herbicide

**Glywell-71** is non-selective, Post emergence systemic herbicide

**Crop :** Tea & non crop

**Applied on :** Annual, Perennial, Broad Leaf & Grassy Weeds.

| Chemical Composition :   |                     |
|--|---------------------|
| Ammonium Salt of Glyphosate a.i. (in-situ)<br>(Based on 100% purity) | 71.000% w/w         |
| Ammonium sulphate  | Q.S.                |
| Inert Polyoxyethylene  | 12.500% w/w         |
| Amine Surfactant   | 1.000% w/w          |
| Moisture   | 1.000% w/w          |
| <b>Total</b>   | <b>100.000% w/w</b> |

## HERBICIDE



### PARAMAX

Paraquat Dichloride 24% SL      Contact Herbicide

**PARAMAX** is a non selective, contact, post emergent herbicide.

**Crop:** Potato, Cotton, Rubber, Wheat, Rice, Apple, Grapes

**Dosage Per Ha\*:** 1.25-3 litre per HA in 500 - 700 litre water

**Packing :** 1 Ltr, 500 ml.

*\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.*

Herbicide

### FITCHEM

Pretilachlor 50% EC

**FITCHEM** is a selective pre emergence broad spectrum herbicide.

**Crop :** Generally for Transplanted Paddy

**Applied on :** Panicum repens, Echinochloa crusgalli, Eclipta Alba, etc

**Dosage Per Ha\*:** 1-1.5 litre per HA in 500 - 700 litre water

*\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.*



### FITCHEM SUPER

Pretilachlor 37% EW

Herbicide

**Fitchem Super** is a selective pre-emergence herbicide in transplanted rice  
**Crop :** Echinochloa colonum, E. crusgalli, Cyperus difformis, C. iria, Digitaria sanguinalis, Frimbristylis miliacea, Eclipta alba, Ludwigia parviflora, Monochoria vaginalis

**Packing :** 3 Litre, 600 ML

**Dosage Per HA\* :** 1500-1875 ML per HA in 250-500 litre Water.

*\*These Dosage are only Indicative. Exact Dose may vary according to Crop stage and Intensity of Pest infestation.*



## HERBICIDE



### HIROSHI

Clodinafop-Propargyl 15% WP

Selective Herbicide

**Hiroshi** is a post emergence systemic grass herbicide and is non phytotoxic to winter wheat.

**Crop:** Generally Wheat

**Applied on:** Phalaris minor etc

**Dosage Per Ha\*:** 160 gm per Acre

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

2,4-D Amine Salt 58% SL

Herbicide

**Azura-58** is non selective, contact, post emergent herbicide.

**Crop:** Potato, Cotton, Rubber, Wheat, Rice, Apple, Grapes

**Packing:** 1 Litre, 400 ML

**Dosage Per HA\*:** 1.25-3 Litre per HA in 500-700 litre Water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### AZURA-58



### AZURA-38

2,4-D Ethyl Ester 38% EC

Systemic Herbicide

Azura-38 is used on a wide variety of terrestrial and aquatic broadleaf weeds. It has little effect on grasses. It is a safe and wide-spectrum weedicide for the control of broad-leaved weeds in many crops. It is a selective, systemic weedicide of the Phenoxy acetic group.

**Crop:** Wheat, Sugarcane, Paddy, Maize.

**Applied on:** Annual and perennial weeds

**Dosage Per HA\*:** 500-800 ml/ acre

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## PIRATE

Pendimethalin 30% EC

Selective Herbicide

**Pirate** is a new generation eco friendly, pre emergence selective herbicide. Applies to soil before crop emerges and prevent early germination of weeds.

**Crop** : Cotton, Soyabean, Paddy, Wheat, etc

**Applied on**: Phalaris minor, Eleusine Indica

**Packing** : 1 Ltr, 500 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## CHEMI-GOLD

Bispyribac Sodium 10% SC

Systemic Herbicide

**Chemi-Gold** is a new generation broad spectrum post emergent herbicide containing Bispyribac-sodium as an active ingredient. It effectively controls most of weed species infesting rice crop, both in nurseries and main field.

**Crop** : Rice

**Applied on** : Echinochloa crusgalli, Echinochloa colonum, Ischaemum rugosum, Cyperus difformis, Cyperus iria, Fimbristylis milliacea, Eclipta alba, Ludwigia etc.

**Packing** : 500 ml, 250 ml, 100 ml, 20 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## HITLER

Metribuzin 70% WP

Herbicide

**Hitler** is Pre & Post emergence selective herbicide.

**Crop** : Sugarcane, Potato, Tomato, Wheat, Soyabean, etc.

**Applied on** : Echinochloa spp, Euphorbia Spp, Portulaca spp, etc

**Packing** : 100 gm.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## CARMAAN

Carbendazim 12% + Mancozeb 63% WP Systemic & Contact Fungicide

**Carmaan** is a broad spectrum systemic and contact Fungicide with Strong and curative action.

**Crop** : Cotton, Paddy, Tomato, Mango, Chilli,

**Applied on** : For Blast of Paddy crop, Brown leaf spot.

**Dosage Per Ha\***: 750 gm in 750 litre water

**Packing** : 1 kg, 500 gm, 250 gm.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Carbendazim 50% WP

Systemic Fungicide

**Caviat-50** is a broad spectrum systemic fungicide which effectively controls the diseases of crops. It is used both as curative and preventive for control of diseases in field crops and vegetables.

**Crop** : Paddy, Wheat, Groundnut, Pea, Cotton, Brinjal, Apple, Grapes, Barley.

**Fungal Diseases** : Blast, Stem Rot, False Smut, Tikka Disease, Root Rot, Collar Rot, Anthracnose, Leaf Spot, Powdery Mildew, Anthracnose Scab

**Packing** : 500 gm.



## CAVIAT-50



## MAAN M-45

Mancozeb 75% WP

Contact Fungicide

**Maan M-45** is a Systemic and contact Fungicide with Strong and curative action.

**Crop** : Chillies, Cucurbits, Wheat, Maize, Paddy, etc.

**Fungal Diseases** : Fruitspot, Leafspot, Dampingoff, Rootrot

**Dosage Per Ha\***: 1.5-2 Kg per HA in 750 litre water

**Packing** : 1 kg, 500 gm.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## FUNGICIDE



### SUPER-80

Sulphur 80% WDG

Contact Fungicide

**Super-80** is a Broad spectrum systemic Fungicide with Protective and Curative actions.

**Crop** : Tobacco, Pulses, Vegetable, Cotton, Grapes, etc

**Fungal Diseases** : Downey mildew, leaf blight, damping off, White Rust, etc

**Dosage Per Ha\***: 250-500 gm per HA in 600 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

### SUPER CAP

Fungicide

Captan 50% WP

**Gripper** is a selective post emergence herbicide with broad spectrum activity. Used for one shot control of wide range of broad leaf weeds.

**Crop**: Generally for Wheat

**Applied on**: Melilotus Indica, Melilotus alba, Vicia Satia, etc

**Dosage Per Ha\***: 8 gm per Acre

**Packing** : 8 gm + 200 ml Surfactant (1 Unit)

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### CHEMICOP

Copper Oxychloride 50% WP

Contact Fungicide

**Azura-58** is non selective, contact, post emergent herbicide.

**Crop** : Potato, Cotton, Rubber, Wheat, Rice, Apple, Grapes

**Packing** : 1 Litre, 400 ML

**Dosage Per HA\*** : 1.25-3 Litre per HA in 500-700 litre Water.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## FUNGICIDE



## CONTURO PLUS

Hexaconazole 5% SC

Systemic Fungicide

**Conturo Plus** is a systemic Fungicide of Triazole group with fast curing and protecting actions.

**Crop :** Paddy, Mango and Grapes

**Fungal Diseases :** Sheath blight, brown spot, Powdery Mildew, etc

**Packing :** 1 Ltr., 500 ml, 250ml.r

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Isoprotiolane 40% EC

## HINATA

Herbicide

**Hinata** belongs to dithiolane group. It is a systemic fungicide with curative and protective action. The chemical is absorbed by the leaves and roots and translocate acropetally and basipetally.

**Crop :** Paddy.

**Fungal Diseases :** Blast Of Paddy

**Packing :** 1 Ltr, 500 ml, 250 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## MINATO

Gibberellic Acid 0.001% L

Plant Growth Regulator

**German Gold** is highly effective Plant growth regulator which acts synergistically with Plant metabolism and accelerates the growth functions of the Plant.

**Applied on :** Cotton, Paddy, Sugarcane, Banana, Onion & Groundnut

**Packing :** 1 Ltr, 500 ml, 250 ml, 100 ml.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

## PGR



## CUTTER

Paclobutrazole 23% SC Plant Growth Regulator

This Plant Growth Regulator is Used For Mango , Onion , Carrot . Garlic , Groundnut , Cashew Nut , Potato , Soybean , Black gram , Peas , Green Gram , Chilly , Tomato , Cauliflower , Sponge Gourd , Brinjal & All Other Major Vegetable Crops.

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

Ethephon 39% SL

Plant Growth Regulator

**ED-Pon** is plant growth regulator with systemic properties. It penetrates into the plant tissues, and is translocated and progressively decomposed to ethylene, which positively affects the growth process.

**Crop :** Mango, Pineapple, Coffee, Tomato, Groundnuts, Rubber

**Applied on :** Fruit Ripening, Breaking Alternate Bearing, Flower Induction etc.

**Dosage Per Ha\*:** 800 - 850 ML / Acre

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



## ED-PON



## TAICHI

Triacontanol EW 0.1% Min. Plant Growth Regulator

**Taichi** for increasing the yield of grains, the dry matter content, height of plants, earlier and stronger tillering, longer and better spread of roots,

**Crop :** Cotton, Tomato, Chilli, Rice, Groundnut, Potato

**Dosage Per Ha\*:** 25 to 30 ml/pump (15 lit), 250 - 300 ml/acre

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.

PGR



## BHOOMI-G

Triacantanol 0.05% GR Min Plant Growth Regulator

**Bhoomi-G** is a Plant Growth Regulator & is recommended to increase the yield of crops like Cotton, Chillies, Rice, Tomato, Groundnut and Potato.

Dosage : 25 Kg per Ha

\*These Doses are only Indicative. Exact Dose may vary according to Crop

Triacantanol 0.05% EC

## SICULAN

Plant Growth Regulator

**Siculan** is a selective pre emergence broad spectrum herbicide.

**Crop :** Generally for Transplanted Paddy

**Applied on :** Panicum repens, Echinochloa crusgalli, Eclipta Alba, etc

**Dosage Per Ha\*:** 1-1.5 litre per HA in 500 - 700 litre water

\*These Doses are only Indicative. Exact Dose may vary according to Crop Stage and Intensity of Pest infestation.



### Our Up Comming Products

## FEEERATA

Chlorantraniliprole 0.4% W/W GR  
Insecticide

## DUNZO

Chlorantraniliprole 18.5% W/W GR  
Insecticide





## **SHIVCHEM AGRO PVT. LTD**

**AN ISO 9001 : 2015 CERTIFIED COMPANY**

Plot No. : GL-42, Hall No. 8, Ganptati Dham, Industrial Area Bahadurgarh (Hr)

Customer Care No. :08368755208 Email : shivchemagro@gmail.com